

Gravitational lensing in galaxy redshift surveys

Daniel J. Mortlock & Rachel L. Webster

*School of Physics, University of Melbourne, Parkville, Vic. 3052,
Australia*

1. Introduction

Gravitationally-lensed quasars are both valuable (individually and statistically) and rare. They are mostly discovered through time-consuming re-imaging of known quasars, but can also be found spectroscopically in galaxy redshift surveys. Q 2237+0305 (Huchra et al. 1985) was found in this manner, but remains the only such discovery. Kochanek (1992) predicted that ~ 1 in 10^5 redshifts would yield a lens although this is somewhat pessimistic, as described below. It is also important that lenses discovered in galaxy surveys tend to have bright, low redshift lens galaxies. Such lenses are particularly useful, as shown by the successful measurement of the source-size and transverse velocity in the Q 2237+0305 system (e.g., Wambsganss et al. 1990; Wyithe et al. 1999).

2. Calculation

The calculation, detailed in Mortlock & Webster (1999), of the expected number of lenses, N_{lens} , is based on the formalism developed in Kochanek (1992). The lensing probability of a single galaxy is integrated over the survey population, subject to a number of selection effects. A lens of total magnitude m , in which the quasar images have magnitude m_q and the galaxy has magnitude m_g , is detectable in a galaxy redshift survey if: the quasar-galaxy composite image is brighter than the survey limit (i.e., $m \leq m_{\text{lim}}$); the quasar images are bright enough to be detected in the composite spectrum (which can be quantified by requiring that $m_q - m_g \leq \Delta m_{\text{qg}} \simeq 2$; Kochanek 1992); and the galaxy is bright enough for the composite object to be classified as non-stellar, and hence enter the redshift survey at all (which can be characterised by the condition that $m_g - m_q \leq \Delta m_{\text{gq}} \simeq 0$). The inclusion of the quasar light in the calculation of m is critical to this calculation, as the sensitivity to lenses is increased by ~ 1 mag, and the number of lenses is increased by up to an order of magnitude, relative to the results of Kochanek (1992).

3. Results

Figure 1 shows the number of lenses expected per 10^4 redshifts as a function of survey depth. Several values of Δm_{qg} are shown, as this represents the largest single uncertainty in the calculation. Note that, at least at the depths of the current redshift surveys, there is very little dependence on the cosmological model, which stands in marked contrast to the usual quasar lensing probability.




Figure 1. The expected number of lenses, N_{lens} , per 10^4 measured redshifts in a galaxy redshift survey of magnitude limit m_{lim} . The three sets of lines are for different values of the “spectral prominence” parameter, Δm_{qg} , as indicated. In each case results are shown for three cosmological models: Einstein-de Sitter (solid lines); empty (dashed lines); and empty and flat (dot-dashed lines).

These results imply that the a posteriori likelihood of a lens being discovered in the Center for Astrophysics (CfA) survey is ~ 0.03 , which is much greater than the previous estimate of ~ 0.003 (Kochanek 1992).

The new generation of surveys – in particular the Sloan Digital Sky Survey (SDSS) and the 2 degree Field (2dF) redshift survey – will dwarf the CfA survey, and should result in the discovery of many new lenses. The SDSS will obtain $\sim 10^6$ high quality galaxy spectra (i.e., $\Delta m_{\text{qg}} \gtrsim 4$) to $m_{\text{lim}} \simeq 19$, which implies $N_{\text{lens}} \simeq 100$. The 2dF redshift survey will contain a quarter the number of galaxies, but has $m_{\text{lim}} = 19.5$, and should yield ~ 10 lenses. Most of these lenses will have deflector redshifts of $\lesssim 0.2$, and in several cases the observer-deflector distance should be comparable to that in the Q 2237+0305 system.

References

- Huchra, J.P., Gorenstien, M., Kent, S., Shapiro, I., Smith, G., Horine, E., & Perley, R. 1985, *AJ*, 90, 691
Kochanek, C.S. 1992, *ApJ*, 397, 381
Mortlock, D.J., & Webster, R.L. 1999, *MNRAS*, in preparation
Wambsganss, J., Paczyński, B., & Schneider, P., 1990, *ApJ*, 358, L33
Wyithe, J.S.B., Webster, R.L., Turner, E.L., & Mortlock, D.J., 1999, *MNRAS*, submitted